

Identifyn Standards for Optical Microscopy Methods™Identifyn® Primary Antibody**Rabbit anti-Human RAD50 Recombinant Monoclonal, IgG - (BLR026N)***Cat ID# - PA-RR 000011**Expected MW: 154 kDa*

HeLa cells were treated with 2 μ M etoposide for 20 hours to induce DNA damage. Cells were homogenized using an ultrasonication probe and a nuclear lysis buffer, then centrifuged to pellet the cell debris. The lysate supernatant (100 μ g) was mixed with [NuPAGE™ LDS Sample Buffer \(4X\)](#), heated at 80 °C for 10 min, and subjected to electrophoresis on [NuPAGE™ Tris-Acetate Mini Protein Gels \(3-8%, 1.0 mm\)](#) in a [Mini Gel Tank](#). Thermo Fisher™ [HiMark™ Pre-stained Protein Standard](#) was used as a molecular weight marker.

Post-electrophoresis, the proteins were transferred to a polyvinylidene difluoride (PVDF) membrane using the [iBlot™ 3 Starter Kit PVDF](#). The transfer membrane was blocked with [Blocker™ Casein](#) for 1 hour, then washed 5 times with 1X PBS. The blocked membrane was incubated with Rabbit anti-Human RAD50 Recombinant Monoclonal, IgG - (BLR026N) for 20 hours, and then washed 5 times in 1X PBS. Identifyn™ SRM Alexa Fluor™ 647 Affinity Purified Donkey Anti-Rabbit, IgG (H + L) was incubated for 1 hour, followed by 5 washes in 1X PBS. The membrane was dried for 10 minutes at room temperature before imaging.

Scanner = [Azure Sapphire FL](#) with Customizable Laser and Filter Modules, Enabling UV-NIR and Z-Plane Imaging.

Scan Settings

Detection mode = Fluorescence

Laser = 638 nm

Emission Filter = 676 $\pm$ 37 nm

Detector = PMT

Intensity = L3

Pixel size = 100 μ m

Scan Speed = High

Quality = High

Focus = Membrane (0 μ m in Z-axis)**Post Processing**

NONE

Image: S. Khanal

© Copyright 2026 GMD12 LLC.